


Bromeli Advisory

DECEMBER 2017

In This Issue

President's Message	2
Post-Mortem on Hurricane Irma	2-5
Upcoming Events	5
Messages	5
Membership Renewal Form	6



BromeliAdvisory

Stop and Smell the Bromeliads

December 2017

WEBPAGE: <http://www.bssf-miami.org/>



<http://www.facebook.com/groups/BromeliadSF/?bookmark t=group>



<http://www.facebook.com/pages/Bromeliad-Society-of-South-Florida/84661684279>

Alex Bello, President
Sandy Roth, VP
Fred Sussenberger, Treasurer
Lenny Goldstein, Secretary
Robert Meyer, Editor

DIRECTORS

Past Pres.: Sandy Roth

Directors:

Barbara Sparling '16-'17
Karen Bradley '17-'18
Ellis Brown '17-'18
Stephanie LaRusso '17-'18

Advertising: Robert Meyer
Door Prize: Alan Herndon
Education: Alan Herndon
Hospitality: Elaine Mills
Library: Barbara Partagas
Membership: Maureen Adelman/Melody Ray
Mem. Plant Sales: Alex Bello
Raffle Table : Melissa Brail
Refreshments: Sandy Roth

What	Who
Sales Table	N/A

DECEMBER 19, 2017 AT 7:00 PM AT FAIRCHILD CORBIN A OR AUDITORIUM
SPEAKER: NONE
RAFFLE TABLE: NONE
FOOD: You – **Holiday Party!!!!!!!!!!!!!!!!!!!!!!**

Smile and Be Ready for the Holiday Party



President's Message

by Alex Bello



Happy Holidays from the Board

Post-Mortem on Hurricane Irma in Southeast Florida

by Leonard Goldstein

Those of us who settled in the southeastern counties of mainland Florida after 1960 have experienced very few hurricanes, but that doesn't mean they weren't memorable.

In 1965, Hurricane Betsy made a shocking clockwise pirouette while already well north of the Bahamas, some 350 miles east of Daytona Beach, and then headed southwestward. It eventually hit Key Largo as a Category 3 hurricane. Flooding was widespread in Broward and Dade counties, and Key Biscayne was reported to have been left almost completely underwater. The storm made a second landfall in Grand Isle, Louisiana.

There wasn't another strong hurricane in southeastern Florida for the next 27 years. Then Andrew devastated the southern third of Miami-Dade County with sustained winds that were upgraded years later to Category 5 status. Both trees and homes were demolished, and The Miami Herald called it 'The Big One.' But the storm was very compact, and areas from Coral Gables north went relatively unscathed, so for two-thirds of the county, Andrew was not The Big One.

Since Andrew in 1992, our part of the state has had a few near-misses, two of which could have caused the most catastrophic hurricane damage ever recorded in U.S. history. The first was Ivan in 2004. The most ominous projection by the National Hurricane Center had Ivan merging from westcentral Cuba as a Category 5 storm on a track due north along the western borders of Miami-Dade, Broward and Palm Beach counties. That meant that the three most populous counties in the state would have been in the 'dirty' northeast quadrant of the hurricane. Fortunately, by the time the midmorning advisory was released a few hours later, the storm track had shifted west into the Gulf of Mexico, and we were spared.

Hurricane Irma was the second near-catastrophe. Its 11th-hour westward track shift saved mainland southeastern Florida from a crippling hit. Most homes experienced little or no wind damage, unless, of course, something fell on them. But lots of vegetation in our area suffered a different fate; broken and overturned trees littered the regional landscape. In light of that significant destruction, many of us were startled when measurements revealed that our sustained winds did not exceed tropical storm force; hurricane-force gusts occurred only periodically.

The positive news to report in the wake of Irma is that palms did very well in mainland southeastern Florida. There are at least a couple of likely explanations. First, the architecture of palms makes them well-suited to withstand strong winds. Fronds of single-trunked pinnate genera such as *Roystonea*, *Chambeyronia*, and *Veitchia*, which carry a small number of leaves, tended to snap off to one degree or other, allowing the plant to present a relatively slender profile against storm winds. Even species with leafier crowns, such as *Satakentia* and *Phoenix roebelenii*, withstood Irma well.

Palmate species also held up admirably, but with less consistent results. Those species which are characterized by a dense crown of large, stiff leaves presented a broader profile against the winds.



For instance, occasional *Bismarckias* went down. Although the usual admonition not to prune trees just in advance of a tropical storm is intended to address the risks posed by unsecured trimmings, it might be worthwhile to make an exception for densely-crowned palms. Pruned fronds could be stashed in a carport or garage for the duration of the storm.

Second, many popular species are native to cyclone-prone regions, and are thus adapted to withstand strong winds. *Coccothrinax*, *Thrinax*, *Sabal* and *Copernicia* were among the numerous palm species which tended to remain upright through Irma while many broadleaf trees were falling all around them.

Just why did those trees perform so poorly in a 'mere' tropical storm? There's a bit of mystery involved in assessing exactly what happened. Can you guess the source of this language?: "Many trees will be snapped or uprooted, blocking numerous roads." It's the Saffir-Simpson Hurricane Wind Scale's description of a Category 3 hurricane, which contains sustained winds of 111-129 mph.¹ Yet it's clearly a pretty accurate description of what Irma did to us.² How does that square with the official claim that our sustained winds were no better than tropical storm-force?

The answer appears to lie in a factor that the Saffir-Simpson Scale doesn't account for: the duration of strong winds in a given area. The consensus among local observers is that what made Irma so destructive of broadleaf trees was the length of time that its winds battered mainland southeastern Florida, rather than their absolute velocity. Hurricane Andrew, for all its power, was fast-moving and relatively dry. Irma, on the other hand, took much longer to push through our area, and produced greater rainfall. After beginning to pound the region on Saturday, it remained capable of unleashing occasional strong gusts as late as sundown on Sunday. Perhaps the Saffir-Simpson Scale could use a little fine-tuning.

A boxing match appears to serve as an apt metaphor for what happened to our broadleaf trees: Imagine a fighter whose opponent administers shots to his midsection for nine rounds. They're not enough to knock him down, but they weaken him, and a solid punch to the chin in the tenth round finally drops him to the canvas. In a similar fashion, Irma, a relatively tame storm locally, softened up many of our trees before delivering the coup de grâce.

What factors made certain broadleaf trees more vulnerable to Irma's extended assault? The explanation generally can be found in the interrelated areas of shape, pruning practices, root system habits, and site preparation. (Also keep in mind that other variables, such as tornadoes and downbursts, within a storm can significantly influence what happens to any tree in your yard, no matter how resistant it might normally be.)

(1) Shape: Dense, broad crowns act like barriers or sails that ultimately prove no match for strong, persistent winds. *Ficus benjamina* is probably Exhibit A in support of this proposition. Its aerial roots, upon contacting soil, form new trunks that help stabilize the tree. When, as is common in our region, those roots are cut above ground, stability is diminished. On the other hand, the root systems of *Ficus benjamina* are extensive and can heave sidewalks, foundations and even interior floor surfaces, so this species poses multiple risks. In recent years, because of high maintenance demands, many of them lining the roads of Coral Gables have been replaced with live oaks, and the differences in storm survival are striking.

(2) Pruning techniques: The practice of hatracking trees only results in a canopy that is denser and thus more wind-vulnerable than before; it has slowly fallen into disrepute and is even illegal in some municipalities. A superior alternative is the technique known as selective pruning, or pickpruning, in which whole branches are removed from the tree, allowing strong winds to flow more freely through the canopy. Selective pruning done in phases over a three-year period appears to achieve the best results. That's particularly true for fruit trees, because reducing one-third of the canopy at a time makes it more likely that they will stay in production throughout the pruning process.

Not all broadleaf species are amenable to even the best of pruning practices. The growth habits of some *Ficus* species and some *Tabebuia* species, for example, involve the interlacing of so many branches that storm-resistant pruning is hard to achieve. In contrast, live oaks are much easier to prune into a very sculptural form in which they develop extremely strong scaffold branches.

(3) Inflexibility: Some tree species are brittle, tending to break up before they fall over. They include avocado, mahogany, royal poinciana, eucalyptus, jacaranda, and silk oak. That doesn't mean that they shouldn't be planted in southern Florida; rather, care should be taken not to place them close to a home, garage, driveway or storage shed. Two other brittle species, bischofia and earleaf acacia, have other strikes against them that render them undesirable for our area.



Fortunately, there are many species that possess greater resilience to tropical storm winds. In addition to live oak, they include lychee, sapodilla, mango, gumbo limbo, mastic, Simpson stopper, strangler fig, cordia, and pandanus. Less well-known than these species is *Elaeocarpus decipiens*, the Japanese Blueberry, but it merits more extensive use. It grows naturally in an attractive pyramidal shape, but takes well to pruning

(4) Root system habits: Not only are some tree species naturally shallow-rooted, but in parts of southeastern Florida that trait is exacerbated by the limestone substrate that occurs just under the surface of the soil. Avoiding species that are notoriously shallow-rooted, or at least exiling them to distant parts of your property, will make your home more likely to escape devastation from tropical storms.

(5) Site preparation: This factor is at the core of many tree failures in our region. In areas where thin soil is underlain by limestone, the practice of planting a tree in a hole just large enough to accommodate its root system is as ill-considered as it is common. The roots of many tropical hardwoods have difficulty pushing their way through the rock. As a result, root systems fail to achieve the type of distribution that leads to optimum stabilization of the tree. A trip down U.S. 1 in southern Miami-Dade County presented stark visuals of that phenomenon. Most of the overturned ornamental trees in the median or along the fringes of the highway exposed a rootball that was small in comparison to the size of the tree. (Perhaps the worst example of lazy planting habits appeared on one large upturned tree, where the side of its container still clung to the rootball! Note to contractors: Remove trees from their pots before planting.)

Although digging a large planting hole is advisable, filling that hole with rich soil is not. That's true whether the site consists of limestone or sand. The practice only encourages the tree's root system to confine itself to the better soil, ultimately affecting the size and stability of the tree. The superior practice is to refill the planting hole with the material that came out of it -- except for any large chunks of rock that may still be present. That technique will encourage the root system to reach out in all directions around the planting hole. There is an additional helpful step in site preparation that can be taken, especially if a backhoe is available. An X-shaped trench can be dug, providing even more room for stabilizing roots to extend into as the tree grows.

1 The NOAA team that updated the Saffir-Simpson Scale in 2012 wrote, "The maximum sustained [emphasis added] surface wind speed (peak 1-minute wind at the standard meteorological observation height of 10 m [33 ft] over unobstructed exposure) associated with the cyclone is the determining factor in the scale."

2 The harm to vegetation that Irma caused in mainland southeastern Florida would also have fit within an earlier Saffir-Simpson Scale description of the effects of a Category 3 hurricane: "Damage to shrubbery and trees with foliage blown off trees and large trees blown down."

Upcoming Events

December 19, 2017
Bromeliad Holiday Party
10901 Old Cutler Road
Coral Gables, FL 33156
Phone: 305-667-1651



January 19-21, 2018

Chocolate Festival

10901 Old Cutler Road
Coral Gables, FL 33156

Phone: 305-667-1651

<http://www.fairchildgarden.org/events-community-outreach/events-details/artmid/486/articleid/1164/12th-annual-international-chocolate-festival>



April 21-22, 2018

Bromeliad Annual Show and Sale

10901 Old Cutler Road Coral Gables, FL 33156
Phone: 305-667-1651

World Conference

May 29, 2018-June 3, 2018
San Diego, CA



If You Got'em, Donate'em Crystal or Give Donations for Awards

Anyone who has previously well deserved and won crystal from the Bromeliad Show is urged to recycle and give the same to Barbara Sparling who handles the awards for the show.

Anyone wanting to donate for awards needs to merely give Barbara an email or call.

Time to Renew Your Membership

Attached is a renewal of membership form. If not received by early next year, you will **not** be included in the membership book and receive other benefits. Thanks for your appreciation.

Badges Wear Them

We make some pretty neat badges. They help our fogged aged memories for names. Starting next month, anyone who wants a door prize **must** be wearing a name badge. This way we will better address one another by name. FYI – do not put the badge onto anything which may be affected by an external magnet.

MESSAGES

NEW MEMBERS PLEASE COLLECT YOUR BADGES

We have 16 new members. Your membership entitles you to Membership Badges. We have them. Come to our meeting and pick them up. Ask for Maureen. She holds the badges.



**BROMELIAD SOCIETY OF SOUTH FLORIDA
BSSF, Inc.
2018 MEMBERSHIP RENEWAL**

(PLEASE PRINT) NAME(S) _____

(Couples - please include BOTH first names)

ADDRESS: _____

CITY _____ STATE _____ ZIP _____

TELEPHONE _____

ADD'L. TEL. _____

Year Joined BSSF: _____

Referred by: _____

Include e-mail address in the roster? (Y/N) ☐ Yes ☐ No

E-mail BromeliAdvisory? (Y/N) ☐ Yes ☐ No (You will not receive a hard copy)

EMAIL _____

IF ANY OF THE ABOVE INFORMATION IS DIFFERENT FROM LAST YEAR,
PLEASE **HIGHLIGHT** OR UNDERLINE

Please renew my(our) membership in the BROMELIAD SOCIETY OF SOUTH FLORIDA
for the period of March 1, 2018 to February 28, 2019.

Check the category: ☐ Single membership \$25.00
☐ Dual membership \$35.00

Dual members enjoy all the privileges of active membership, but receive only one
copy of the BromeliAdvisory.

**DUES MUST BE RECEIVED BY February 28, 2018, IN ORDER FOR YOU TO BE
INCLUDED IN THE 2018 ROSTER**

Please mail completed form and check

(Payable to BSSF, Inc.) to:

Maureen Adelman
9421 SW 134 Street
Miami, FL 33176

Or E-Mail this form to:

mhadelman@comcast.net

You can pay up to 5 years at the rates described above. All payments are
NONREFUNDABLE